

# Colonoscopy Prep –

## GoLyte Split Prep

Please review this entire document! These instructions are your physician's specific instructions. Follow all steps carefully to ensure successful prep and procedure. If you are or may be pregnant, please discuss the risks and benefits of this procedure with your provider. Questions? Call 317.214.5468.

<p><b>10 Days</b> before your procedure</p>	<p><b>Review and start following all MEDICATION GUIDELINES on the <a href="#">here</a>.</b></p> <ul style="list-style-type: none"> <li>• Consult with your prescribing provider before discontinuing any blood thinners/ anticoagulants and/or diabetic medications.</li> </ul>
<p><b>3 Days</b> before your procedure</p>	<p><b>Start following LOW FIBER/LOW RESIDUE DIET on <a href="#">here</a>.</b></p> <ul style="list-style-type: none"> <li>• Stop eating nuts, seeds, corn, popcorn, raw fruits, raw vegetables, whole wheat foods, multi-grain foods, bran and fiber supplements until after your procedure.</li> <li>• Stop taking all vitamins &amp; supplements.</li> </ul>
<p><b>2 Days</b> before your procedure</p>	<ul style="list-style-type: none"> <li>• Continue following LOW FIBER/LOW RESIDUE DIET.</li> <li>• Take medications as usual unless directed otherwise.</li> <li>• CONFIRM arrangements with your friend/family member adult driver (age 18+).</li> <li>• <b>DO NOT eat solid foods after midnight.</b></li> </ul>
<p><b>1 Day</b> before your procedure</p>	<p><b>Start following CLEAR LIQUID DIET included in this document.</b></p> <ul style="list-style-type: none"> <li>• NO SOLID FOODS, CLEAR LIQUIDS ONLY</li> <li>• Drink plenty of clear liquids throughout the day to avoid dehydration.</li> </ul> <p><b>BEFORE 9AM:</b> Add water to the fill line of the GoLyte®, CoLyte®, NuLyte®, GaviLyte® container, mix and refrigerate.</p> <p><b>AT Noon: Take three 5mg tablets of bisacodyl (Dulcolax) available over the counter</b></p> <p><b>AT 5PM:</b> Drink 64oz (half) of prep solution - One 8oz glass every 15 minutes (x8); Refrigerate remaining 64oz of prep solution.</p> <p><b>Take three Simethicone (GAS-X) 125mg tablets available over the counter</b></p> <p><b>IMPORTANT:</b> 2nd dose will be 6 hours prior to appointment arrival time day of procedure.</p>
<p><b>The Day</b> of your procedure</p>	<p><b>6 HOURS PRIOR TO APPOINTMENT ARRIVAL TIME:</b></p> <p>Drink remaining 64oz of prep solution, one 8oz glass every 15 min. (x8). Continue following CLEAR LIQUID DIET until 4 hours prior to appointment arrival time</p> <p><b>4 HOURS PRIOR TO APPOINTMENT ARRIVAL TIME:</b></p> <ul style="list-style-type: none"> <li>• STOP DRINKING, NOTHING BY MOUTH</li> <li>• No eating, drinking, smoking, or vaping. No gum or hard candy.</li> <li>• Take any approved medication with a small sip of water at least 4 hours before procedure</li> <li>• You may brush your teeth, but do not swallow.</li> <li>• <b>You are ready for the colonoscopy if you follow all instructions, and your stool is clear enough to see the bottom of the toilet bowl after a bowel movement.</b></li> </ul>

# Colonoscopy Prep – GoLytely Split Prep

## BEFORE THE PROCEDURE

- If you are on blood thinners/anticoagulants, please contact your prescribing physician before discontinuing your medication. Medication guidelines are included [here](#).
- Colonoscopy prep will likely require the purchase of some over-the-counter medications and food/drinks allowed on the low fiber and clear liquid diets. Shopping list, allowed foods and foods not allowed are included in this document.
- If you are being sedated, please plan to have a friend/family member adult driver (age 18+) for after your procedure.

## THE PREP

- Be sure to have access to a restroom. Individual response time to prep may vary. You may have loose stools.
- You may be uncomfortable/bloated and feel chilled until you start having bowel movements (2-4 hours). Drinking warm, clear liquids such as tea, warm apple juice or hot water can help.
- Alcohol-free baby wipes may help ease skin irritation.
- Use over-the-counter hemorrhoid pads if necessary.

## THE APPOINTMENT

- Bring to the appointment:
  - List of all current medications (including allergies or over the counter)
  - Inhalers
  - Photo ID and up-to-date insurance information
  - Leave valuables (jewelry, smart watches, cell phones) at home or with your driver
- Arrive 1 hour prior to scheduled procedure time.
- You **MUST** have a friend/family member adult driver (age 18+) remain in the building throughout the entire procedure and available to drive you home after your procedure.
- Taxi/cab, Uber/Lyft, bus and medical transport are NOT acceptable rides home unless accompanied by an adult (age 18+) friend/family member.

## **AFTER THE PROCEDURE**

Because of the sedation given during the procedure, you may experience sleepiness, dizziness, forgetfulness or lightheadedness. Your judgement and reflexes may be impaired. These are normal reactions to the medication that can last for the remainder of the day.

## **FOR THE REMAINDER OF YOUR PROCEDURE DAY**

- **DO NOT** drive a vehicle or operate complex machinery
- **DO NOT** perform any strenuous activity
- **DO NOT** drink alcohol
- **DO NOT** smoke unsupervised
- **DO NOT** make any legal decisions
- Additional information will be included on your discharge instructions

## **SHOPPING LIST**

### **Required Items**

- Allowed food for low fiber/low residue diet
- Allowed beverages for clear liquid diet
- GoLytely®, CoLyte®, NuLytely Prep® (polyethylene Glycol)
- 3 bisacodyl 5mg (Dulcolax) tablets available over the counter.
- 3 Simethicone (GAS-X) 125mg tablets available over the counter.



### **Optional Items**

- Hemorrhoidal pads
- Alcohol-free baby wipes

Food Category	Allowed	Not Allowed
<b>Beverages</b>	Coffee, tea, hot chocolate, soda, fruit & vegetable juice without pulp; Less than 2 cups milk/day	No alcohol; No fruit or vegetable juice with pulp; no red or purple
<b>Breads, grains &amp; cereal</b>	White bread, rolls & bagels; Saltines, cheese crackers; Refined dry & cooked cereals including corn flakes, crisp/puffed rice, Cream of Wheat	No whole wheat/whole grain bread, rolls, crackers or cereal; No breads/cereal with bran, oats, seeds, nuts, raisins or dates; No oatmeal
<b>Dairy</b>	Cheese, cottage cheese, plain yogurt, sour cream, buttermilk, cream	No yogurt mixed with granola or berries (or other fruit with skin)
<b>Meat &amp; eggs</b>	Chicken, turkey, fish, seafood, tofu, eggs	No lunch meat, hot dogs, sausage; No spiced/cured meats; No fried, tough, grisly meats
<b>Fats</b>	Butter, cream, mayo, avocado, cooking oils & shortening; creamy peanut/almond butter	No nuts, coconut, spicy salad dressings
<b>Fruits</b>	Ripe, peeled apples, bananas, melon, pears & peaches; cooked/canned fruits without skin/peel/membranes	No raw fruit with seeds, skin, or membranes inc. berries, pineapple, apples, oranges, grapefruit, watermelon, kiwi, pomegranate, dragon fruit
<b>Vegetables</b>	<b>All vegetables must be well-cooked:</b> Asparagus, carrots, mushrooms, pumpkin	<b>Raw vegetables are not allowed</b> No red or golden beets No raw spinach, lettuces or cucumber No corn, peas/snap peas or green/wax beans
<b>Potatoes &amp; Starches</b>	Potatoes without skin (white, yellow, sweet & yams), noodles, white rice, hominy	No purple potatoes, fried potatoes, potato skins, whole wheat pasta, brown & wild rice
<b>Beans &amp; legumes</b>	<b>None allowed</b>	<b>None allowed</b>
<b>Nuts &amp; seeds</b>	<b>None allowed</b>	<b>None allowed</b>
<b>Soups</b>	Cream soups made with allowed milk (less than 2 cups/day) & allowed vegetables, broth soups	Soups with more than 2 cups of milk per serving per day
<b>Miscellaneous</b>	White sauce (made from allowed milk), meat gravy, ketchup, mustard, tomato sauce, chocolate, salt, vinegar, lemon juice, ground spices & herbs in moderate amounts	No garlic, ginger, olives, pickles, popcorn, horseradish, cayenne, chili powder

# Colonoscopy Prep – GoLytely Split Prep

## **NO SOLID FOODS. CLEAR LIQUIDS ONLY.**

Be sure to drink plenty of clear liquids with calories on clear liquid diet days

### **APPROVED CLEAR LIQUIDS**

- Sports drinks/Gatorade®/VitaminWater® No red or purple
- Fruit juice (no pulp) Apple, white grape, white cranberry; No red or purple
- Soda & carbonated beverages; No red or purple
- Clear broth/bouillon Chicken, vegetable, beef flavors
- Gelatin/Jell-O® No red or purple
- Ice pop/Popsicles® No red or purple
- Gummy bears; No red or purple
- Black coffee & tea No liquid or powdered creamer/milk (dairy/soy/nut/oat); Sugar/sweeteners OK
- Water Plain, flavored, carbonated; No red or purple

### **IF YOU ARE DIABETIC, AIM FOR 45 GRAMS OF CARBOHYDRATES PER MEAL AND 15-30 GRAMS PER SNACK**

- **Clear liquids with about 15 grams of carbohydrates**
  - 4oz apple juice, 8oz sports drink, ½ cup gelatin, 2 popsicles/ice pops
  - Clear liquids with zero carbohydrates
  - Black coffee, tea (unsweetened or diet), clear diet soda, seltzer, flavored water, fat-free broth, bouillon or consommé

**⊘ NO RED ⊘ NO PURPLE ⊘ NO DAIRY ⊘ NO PULP**