

# Colonoscopy Prep -

## MiraLAX Split Prep

**Please review this entire document!** These instructions are your physician's specific instructions. Follow all steps carefully to ensure successful prep and procedure. If you are or may be pregnant, please discuss the risks and benefits of this procedure with your provider. **Questions? Call 317-214-5468.**

<p><b>10 Days</b> before your procedure</p>	<p><b>Review and start following all MEDICATION GUIDELINES on <a href="#">here</a>.</b></p> <ul style="list-style-type: none"> <li>Consult with your prescribing provider before discontinuing any blood thinners/ anticoagulants and/or diabetic medications.</li> </ul>
<p><b>1 Day</b> of your procedure</p>	<p><b>Start following CLEAR LIQUID DIET</b> included in this document.</p> <ul style="list-style-type: none"> <li>NO SOLID FOODS, CLEAR LIQUIDS ONLY</li> <li>Drink plenty of clear liquids throughout the day to avoid dehydration.</li> </ul> <p><b>BEFORE 9AM:</b> In a pitcher, add 8.3oz (238g) MiraLAX® powder (polyethylene glycol 3350) in 64oz of Gatorade®. Stir well and refrigerate. Avoid red/purple Gatorade®.</p> <p><b>AT 5PM:</b> Take 3 tablets of Dulcolax® (bisacodyl), total 15mg.</p> <p><b>AT 6PM:</b> Drink 32oz (half) of prep solution - One 8oz glass every 15 minutes (x4); Refrigerate remaining 32oz prep solution.</p> <p>Take three Simethicone (GAS-X) 125mg tablets available over the counter.</p> <p><b>IMPORTANT:</b> 2nd dose will be 6 hours prior to appointment arrival time on day of procedure.</p>
<p><b>The Day</b> of your procedure</p>	<p><b>6 HOURS PRIOR TO APPOINTMENT ARRIVAL TIME:</b> Drink remaining 32oz of MiraLAX®/Gatorade® solution, one 8oz glass every 15 minutes (x4). <b>Continue following CLEAR LIQUID DIET until 4 hours prior to appointment arrival time.</b></p> <p><b>4 HOURS PRIOR TO APPOINTMENT ARRIVAL TIME:</b></p> <ul style="list-style-type: none"> <li>STOP DRINKING, NOTHING BY MOUTH</li> <li>No eating, drinking, smoking, or vaping. No gum or hard candy.</li> <li>Take any approved medication with a small sip of water at least 4 hours before procedure</li> <li>You may brush your teeth, but do not swallow.</li> <li>You are ready for the colonoscopy if you followed all instructions and your stool is no longer formed but clear or yellow liquid (like urine or tea).</li> </ul>

## WHAT TO EXPECT

### BEFORE THE PROCEDURE

- If you are on blood thinners/anticoagulants, please contact your prescribing physician before discontinuing your medication. **Medication guidelines are included [here](#).**
- Colonoscopy prep will likely require the purchase of some over-the-counter medications and food/drinks allowed on the low fiber and clear liquid diets. A shopping list of allowed foods and foods not allowed are included in this document.
- If you are being sedated, please plan to have a friend/family member adult driver (age 18+) for after your procedure.

### THE PREP

- Be sure to have access to a restroom. Individual response time to prep may vary. You may have loose stools.
- You may be uncomfortable/bloated and feel chilled until you start having bowel movements (2-4 hours). Drinking warm, clear liquids such as tea, warm apple juice or hot water can help.
- Alcohol-free baby wipes may help ease skin irritation.
- Use over-the-counter hemorrhoid pads if necessary.

### THE APPOINTMENT

- Bring to the appointment:
  - List of all current medications (including allergy or over the counter)
  - Inhalers
  - Photo ID and up-to-date insurance information
  - Leave valuables (jewelry, smart watches, cell phones) at home or with your driver
- Arrive 1 hour prior to scheduled procedure time.
- You **MUST** have a friend/family member adult driver (age 18+) remain in the building throughout the entire procedure and available to drive you home after your procedure.
- Taxi/cab, Uber/Lyft, bus and medical transport are **NOT** acceptable rides home unless accompanied by an adult (age 18+) friend/family member.

## **AFTER THE PROCEDURE**

Because of the sedation you are given during the procedure, you may experience sleepiness, dizziness, forgetfulness or lightheadedness. Your judgement and reflexes may be impaired. These are normal reactions to the medication that can last for the remainder of the day.

## **FOR THE REMAINDER OF YOUR PROCEDURE DAY**

- **DO NOT** drive a vehicle or operate complex machinery
- **DO NOT** perform any strenuous activity
- **DO NOT** drink alcohol
- **DO NOT** smoke unsupervised
- **DO NOT** make any legal decisions
- Additional information will be included on your discharge instructions

## **SHOPPING LIST**

### **Required Items**

- Allowed food for low fiber/low residue diet
- Allowed beverages for clear liquid diet
- MiraLAX® powder 8.3oz (238g) and Gatorade® 64oz
- Dulcolax® (bisacodyl)
- Three Simethicone (GAS-X) 125mg tablets available over the counter.

### **Optional Items**

- Hemorrhoidal pads
- Alcohol-free baby wipes

## **CLEAR LIQUID DIET**

NO SOLID FOODS. CLEAR LIQUIDS ONLY.

Be sure to drink plenty of clear liquids **with calories** on clear liquid diet days

### **APPROVED CLEAR LIQUIDS**

- **Sports drinks/Gatorade®/VitaminWater®** No red or purple
- **Fruit juice (no pulp)** Apple, white grape, white cranberry; No red or purple
- **Soda & carbonated beverages** No red or purple
- **Clear broth/bouillon** Chicken, vegetable, beef flavors
- **Gelatin/Jell-O®** No red or purple
- **Ice pop/Popsicles®** No red or purple
- **Gummy bears** No red or purple
- **Black coffee & tea** No liquid or powdered creamer/milk (dairy/soy/nut/oat); Sugar/sweeteners OK
- **Water** Plain, flavored, carbonated; No red or purple

**IF YOU ARE DIABETIC, AIM FOR 45 GRAMS OF CARBOHYDRATES PER MEAL AND 15-30 GRAMS PER SNACK**

### **Clear liquids with about 15 grams of carbohydrates**

- 4oz apple juice, 8oz sports drink, ½ cup gelatin, 2 popsicles/ice pops

### **Clear liquids with zero carbohydrates**

- Black coffee, tea (unsweetened or diet), clear diet soda, seltzer, flavored water, fat-free broth, bouillon or consommé

⊘ NO RED ⊘ NO PURPLE ⊘ NO DAIRY ⊘ NO PULP